

AD-A039 520

AEROSPACE CORP EL SEGUNDO CALIF MATERIALS SCIENCES LAB  
CARBON: OCCURRENCE OF LINEAR FORMS IN NATURAL GRAPHITE. (U)  
APR 77 A G WHITTAKER

F/G 20/2

UNCLASSIFIED

TR-0077(2250-10)-2

SAMSO-TR-77-94

F04701-76-C-0077

NL

| OF |

AD  
A039 520



END

DATE  
FILMED

6-77

AD A 039520

(13)  
NW

## Carbon: Occurrence of Linear Forms in Natural Graphite

Materials Sciences Laboratory  
The Ivan A. Getting Laboratories  
The Aerospace Corporation  
El Segundo, Calif. 90245

15 April 1977

APPROVED FOR PUBLIC RELEASE;  
DISTRIBUTION UNLIMITED

DDDC  
MAY 17 1977


Prepared for  
SPACE AND MISSILE SYSTEMS ORGANIZATION  
AIR FORCE SYSTEMS COMMAND  
Los Angeles Air Force Station  
P.O. Box 92960, Worldway Postal Center  
Los Angeles, Calif. 90009


AD No. \_\_\_\_\_  
DDC FILE COPY,

This interim report was submitted by The Aerospace Corporation, El Segundo, CA 90245, under Contract No. F04701-76-C-0077 with the Space and Missile Systems Organization, Deputy for Advanced Space Programs, P.O. Box 92960, Worldway Postal Center, Los Angeles, CA 90009. It was reviewed and approved for The Aerospace Corporation by W. C. Riley, Director, Materials Sciences Laboratory. Lieutenant A. G. Fernandez, SAMSO/YAPT, was the project officer for Advanced Space Programs.


This report has been reviewed by the Information Office (OI) and is releasable to the National Technical Information Service (NTIS). At NTIS, it will be available to the general public, including foreign nations.

This technical report has been reviewed and is approved for publication.

  
Arturo G. Fernandez, Lt, USAF

  
Joseph Gassmann, Major, USAF

FOR THE COMMANDER

  
Floyd R. Stuart, Colonel, USAF  
Deputy for Advanced Space Programs

ADMISSION for	
NTIS	Write Sigin <input checked="" type="checkbox"/>
DDC	Diff Section <input type="checkbox"/>
UNANNOUNCED	<input type="checkbox"/>
JUSTIFICATION.....	
BY.....	
DISTRIBUTION/AVAILABILITY CODES	
DISC.	AVAIL. OR/OF SPECIAL
A	



UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

19 REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM	
18 REPORT NUMBER SAMS0 TR-77-94	2 GOVT ACCESSION NO.	3 RECIPIENT'S CATALOG NUMBER	
6 TITLE (and Subtitle) CARBON: OCCURRENCE OF LINEAR FORMS IN NATURAL GRAPHITE.		5 TYPE OF REPORT & PERIOD COVERED Interim rept.	
7 AUTHOR(s) A. Greenville/Whittaker		6. PERFORMING ORG. REPORT NUMBER TR-0077(2250-10)-2	
9. PERFORMING ORGANIZATION NAME AND ADDRESS The Aerospace Corporation El Segundo, Calif. 90245		8. CONTRACT OR GRANT NUMBER(s) F04701-76-C-0077	
11. CONTROLLING OFFICE NAME AND ADDRESS Space and Missile Systems Organization Air Force Systems Command Los Angeles, Calif. 90009		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) 12 13p.		12. REPORT DATE 15 Apr 77	
		13. NUMBER OF PAGES 11	
		15. SECURITY CLASS. (of this report) Unclassified	
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release; distribution unlimited			
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)			
18. SUPPLEMENTARY NOTES			
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Carbon Layer Structure of Carbon Crystals Linear Carbon Forms Linear Carbon Forms, Age Linear Carbon Forms, Geographical Distribution Natural Carbon			
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) A study of several natural carbon formations, from geographical locations ranging from Celyon to California, revealed that the carbons contain small amounts of linear carbon forms, e.g., lchaoite, α carbyne, and β carbyne. The results indicate that these "rare" carbon forms have worldwide distribution and, as with diamond, are prevented from tranformation to graphite by a high kinetic barrier.  ALPHA      BETA			

DD FORM 1473  
(FACSIMILE)UNCLASSIFIED  
SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

1B

## PREFACE

C. H. Hills helped with the x-ray work, E. S. Watts the electron diffraction, N. Marquez the ion probe, A. L. Palyo the scanning electron microscopy, and M. E. Neudorffer the collection and preparation of the samples. The author gratefully acknowledges these coworkers for their assistance.

## CARBON: OCCURRENCE OF LINEAR FORMS IN NATURAL GRAPHITE

Heretofore the linear carbon forms were believed to be rare forms found only in meteorites or meteorite craters or produced in trace quantities under unusual laboratory conditions. However, a study of natural carbon deposits indicates that one or more of these carbon forms commonly occur over a very wide geographic area.


Since these carbon forms are not well known, background on their properties is necessary. Each crystal form of carbon gives a characteristic negative-ion spectrum under conditions that prevail in an ion microprobe.<sup>1,2</sup> Because these phenomena were used extensively in the study of the linear carbon forms,<sup>3,4,5,6</sup> graphite from various natural sources was studied to check the universality of the method and to improve the accuracy of the graphite reference spectrum.

Another characteristic of the carbon forms was revealed by studying the etch patterns produced by the ion microprobe beam with a scanning electron microscope (SEM). In diamond and graphite (both in the a and c directions), the etch pattern is a smooth, essentially featureless, depression. The linear carbon forms and lonsdaleite etch into a variety of patterns such as pinnacles, sharp steps, and rounded steps. Although the detailed association between the patterns and carbon forms is far from established, the patterns are useful. At present, eight linear forms can be distinguished by their  $a_0$  values. Unfortunately,  $a_0$  and  $c_0$  are known for only chaoite,  $\alpha$  and  $\beta$  carbyne,



and carbon VI and VII. Therefore, positive identification of all the linear forms is not possible at present. A few of these forms are very hard; Chaoite, for example, is harder than  $B_4C$ , which is a superhard material. Others are as soft or softer than graphite. For the hexagonal carbon forms, there is a correlation between the etch pattern and the hardness. The soft forms show smooth shallow depressions; the hard forms show deep holes with the characteristic patterns on the walls and bottom. The hard forms also have a higher sputtering rate.


The first crystals studied were obtained from coarse crystalline marble from a quarry in Ticonderoga, New York. The crystals were removed from the marble with hydrochloric acid and subsequently washed in distilled water. They were analyzed with ARL ion microprobe mass analyzer (IMMA) Model 101000 in both the positive- and negative-ion secondaries from the samples. The positive-ion spectrum indicated very few impurities in trace amounts; in the negative-ion spectrum, the results differed, depending on the location probed. At A in Fig. 1(a), a typical graphite spectrum was obtained. At B, the spectrum is characteristic of the linear carbon forms. The probe holes produced at these locations were examined with an SEM. The hole at A is the typical featureless depression characteristic of graphite and is in agreement with the  $C_n^-$  ion spectrum obtained at that point. The probe hole produced at B had a complex etch pattern, Fig. 1(b). In this case, the ion beam just grazed the vertical face of a growth step on the crystal surface. It is evident that the crystal is not a single form but consists of layers of different carbon forms. The first layer is a very thin layer of a linear form.




(a)



(b)



(c)



(d)

Fig. 1. Carbon Crystals. (a) Carbon crystal from Ticonderoga marble; (b) Same crystal as in (a) showing magnified image of ion probe hole at B; (c) Carbon crystal from Santa Rosa Mountains marble showing probe holes on a prism face; (d) Same crystal as in (c) showing mechanically damaged edge.



The second is a thin layer of what appears to be graphite. The third layer is one of the linear carbon forms. The fourth layer may be graphite, as indicated by the smooth etch surface. The fifth, sixth, and seventh are linear forms. The fifth and seventh are of the same form, but the sixth is different. The step edges of the sixth layer are sharper than those of the adjacent layers. More important, it is brighter and, therefore, gives a higher yield of secondary electrons than the form in the adjacent layers. The eighth layer appears to be graphite. Other layers of linear forms appear as the major body of the crystal is penetrated. Although this growth step represents only a small fraction of the total thickness of the crystal, the SEM photograph indicated that there was a high enough concentration of linear forms to produce lines or spots on an x-ray diffraction (XRD) pattern. This prediction was correct, and the d-values obtained are shown in Table 1. The crystal was rotated about an axis nearly perpendicular to the c-axis in order to minimize orientation effects. From these data, it is evident that there are many more reflections than would be obtained from graphite. Since the IMMA showed a very low impurity level, the nongraphite reflections must result from other carbon forms. Indeed, the crystal probably contains very little graphite because two prominent graphite reflections, the 1.23- and 1.16-Å spacings, are absent.

The crystal was mounted and sliced by means of an ultramicrotome in the hope that high-energy electron diffraction (HEED) could be performed at suitable places across the slice, and thereby the layers could be identified unambiguously. Unfortunately, the soft layers were crushed, and the hard

Table 1. X-ray diffraction data from carbon crystals obtained  
from Ticonderoga and Santa Rosa Mountain marble

Ticonderoga		Santa Rosa	
Observed d, Å	Relative <sup>a</sup> Intensity	Observed d, Å	Relative Intensity
6.607	S	3.818	S
4.133	S	3.453	S
3.648	M	3.069	M
3.376	VS	2.659	M
2.122	M	2.132	S
2.014	M	2.032	S
1.793	W	1.804	S
1.675	S	1.549	S
1.544	W	1.321	M
1.519	W	1.290	W
1.315	VVW	1.234	W
1.137	VVW	1.158	S
1.119	M	1.137	M
1.060	VVW	1.055	W
1.047	VW		
1.010	VVW		
0.9906	VW		
0.9584	VW		
0.8739	W		
0.8384	M		

<sup>a</sup> Intensities are graded in the order of increasing intensity  
as follows: VVW - VW - W - M - S - VS

layers crumbled, destroying the relationship between the layers and the diffraction patterns. However, the HEED results overlapped the x-ray data rather well; and, as usual, each technique produced reflections that the other did not. On the basis of the negative ion spectra and the diffraction data, the following tentative assignments were made: the fifth and seventh layers are  $\beta$  carbyne; and the sixth is  $\alpha$  carbyne or chaoite.

Ten crystals from Ticonderoga marble were studied. Three or four of them consisted of graphite only; the rest had a layered structure and linear forms. However, this was a poor sampling of a fairly large geological formation and may not be representative of the entire formation.

A carbon crystal obtained from a graphitic marble from the Santa Rosa Mountains, Imperial County, California, also was analyzed. Again, the positive-ion spectra showed very few impurities in trace amounts, and the negative ion spectrum depended on the location probed. Several probe holes on a prism face (hexagonal crystal) are shown in Fig. 1(c). The layer structure is evident, and the etch patterns indicate that about two-thirds of the crystal thickness is of a soft form and one-third of the thickness is a hard form. This conclusion is supported by the appearance of a mechanically damaged edge, Fig. 1(d). The region showing the smooth etch pattern deformed as a soft material, and in the region showing the deep holes with clear etch patterns there is evidence of brittle fracture.

The x-ray diffraction results are shown in Table 1. Many nongraphite reflections were observed. Note that the strong reflection at  $3.45 \text{ \AA}$  occurs at a d-value that is high for a well-formed graphite crystal and that the



reflection at 1.234 Å is weak, whereas the one at 1.158 Å is strong. In graphite, these reflections are of nearly equal intensity. None of the carbon negative-ion spectra indicated the presence of graphite. Therefore, this crystal contains no graphite, even though a superficial inspection of the XRD data suggests that the crystal is an impure graphite crystal. Another indication that the crystal is not graphite is its shape. Graphite crystals have hexagonal symmetry, whereas this crystal has trigonal symmetry, which is the symmetry seen most often in crystals of the linear carbon forms.

Similar results were obtained for carbon samples obtained from Ceylon; Fossil Canyon; the Coyote Mountains, Imperial County, California; Pacoima Canyon, the San Gabriel Mountains, California; and Coon Creek Canyon, the San Bernardino Mountains, California. Pacoima Canyon material was fractionated according to density. The fraction between 2.9 to 3.3 g/ml was a fairly pure fraction of a form tentatively designated as carbon VII. This study is continuing, and the final detailed results will be published later. At this time, the above findings suggest that the linear carbon forms can be found all over the world. As with diamond, they are thermodynamically unstable and have a kinetic barrier to transformation that allows them to exist for at least 60 million years.

## REFERENCES

1. W. K. Stuckey and A. G. Whittaker, Paper No. TP-177, presented at 10th Carbon Conference, Lehigh University, Bethlehem, Pennsylvania, 27 June to 2 July 1971.
2. Both the Cameca direct imaging mass analyzer and Applied Research Laboratory ion probe gave the same negative ion spectra.
3. A. M. Sladkov and Yu. P. Kudryavtsev, Priroda 5, 37 (1969).
4. A. El Goresy and G. Donnay, Science 161, 363 (1968).
5. A. G. Whittaker and P. L. Kintner, Science 165, 589 (1969);  
A. G. Whittaker and G. M. Wolten, Science 178, 54 (1972).
6. V. I. Kasatochkin, V. V. Savianskiy, G. N. Smirnov, and  
V. M. Mel'nichenko, Doklady, Akademicheskikh Nauk SSR 217, 796 (1974).

#### THE IVAN A. GETTING LABORATORIES

The Laboratory Operations of The Aerospace Corporation is conducting experimental and theoretical investigations necessary for the evaluation and application of scientific advances to new military concepts and systems. Versatility and flexibility have been developed to a high degree by the laboratory personnel in dealing with the many problems encountered in the nation's rapidly developing space and missile systems. Expertise in the latest scientific developments is vital to the accomplishment of tasks related to these problems. The laboratories that contribute to this research are:

Aerophysics Laboratory: Launch and reentry aerodynamics, heat transfer, reentry physics, chemical kinetics, structural mechanics, flight dynamics, atmospheric pollution, and high-power gas lasers.

Chemistry and Physics Laboratory: Atmospheric reactions and atmospheric optics, chemical reactions in polluted atmospheres, chemical reactions of excited species in rocket plumes, chemical thermodynamics, plasma and laser-induced reactions, laser chemistry, propulsion chemistry, space vacuum and radiation effects on materials, lubrication and surface phenomena, photo-sensitive materials and sensors, high precision laser ranging, and the application of physics and chemistry to problems of law enforcement and biomedicine.

Electronics Research Laboratory: Electromagnetic theory, devices, and propagation phenomena, including plasma electromagnetics; quantum electronics, lasers, and electro-optics; communication sciences, applied electronics, semiconducting, superconducting, and crystal device physics, optical and acoustical imaging; atmospheric pollution; millimeter wave and far-infrared technology.

Materials Sciences Laboratory: Development of new materials; metal matrix composites and new forms of carbon; test and evaluation of graphite and ceramics in reentry; spacecraft materials and electronic components in nuclear weapons environment; application of fracture mechanics to stress corrosion and fatigue-induced fractures in structural metals.

Space Sciences Laboratory: Atmospheric and ionospheric physics, radiation from the atmosphere, density and composition of the atmosphere, aurorae and airglow; magnetospheric physics, cosmic rays, generation and propagation of plasma waves in the magnetosphere; solar physics, studies of solar magnetic fields; space astronomy, x-ray astronomy; the effects of nuclear explosions, magnetic storms, and solar activity on the earth's atmosphere, ionosphere, and magnetosphere; the effects of optical, electromagnetic, and particulate radiations in space on space systems.

THE AEROSPACE CORPORATION  
El Segundo, California